

Préime DermaFacial

CONTRAINDICATIONS

Local Contraindications

These apply to specific treatment areas and do not necessarily prevent full-face treatment:

- Active skin infections (e.g., herpes simplex, bacterial or fungal infections)
- Open wounds, cuts, broken skin, or burns
- Active eczema, psoriasis, dermatitis, or rashes
- Severe rosacea or during flare-ups
- Inflamed, pustular, or cystic acne
- Prominent platysma bands (*MicroT only*)
- Thin, fragile, or highly vascular skin (*AquaB & VibroX only*)
- Severe melasma or high risk of post-inflammatory hyperpigmentation (*Collagen+ only*)
- Metal implants, screws, or plates in or near the treatment area (*MicroT, Collagen+ & UltraB only*)
 - Note: Dental implants and orthodontic braces are not contraindicated.

General Contraindications

These apply to all handpieces and should exclude treatment entirely:

- Impaired skin sensation or neuropathy in the treatment area
- Sunburn or severely compromised skin barrier
- Uncontrolled diabetes (due to impaired wound healing)
- Severe autoimmune or collagen disorders (e.g., lupus, scleroderma)
- Recent use of isotretinoin (Roaccutane) — must be discontinued for at least 6 months
- Use of anticoagulant medications (e.g., warfarin, heparin, prescription-strength aspirin) — may increase risk of bruising, bleeding, or delayed healing
- Pregnancy or breastfeeding — treatment requires prior medical clearance
- Recent advanced skin procedures (e.g., microneedling, chemical peels, injectables) — wait 2–6 weeks depending on healing
- Recent Botox or dermal filler injections — wait 2–4 weeks before treating affected area
- Use of retinoids, AHAs, BHAs, or exfoliating acids within the past 48–72 hours
- Presence of pacemakers, defibrillators, or other implanted electronic devices (*MicroT, Collagen+ & UltraB only*)
- Patients under the age of 18 (*MicroT, Collagen+ & UltraB only*)
- Known allergies to product ingredients, aspirin, or fragrance allergens (e.g., limonene, linalool, benzyl benzoate)